UNIVERSAL JOURNAL OF MEDICAL AND NATURAL SCIENCES
ISSN 2992-8826
https://sr-journals.org/index.php/UJMANS

ELECTRICAL CURRENT IN LIVING ORGANISMS: BIOPHYSICAL MECHANISMS AND APPLIED ANALYSIS

Usanova Marjona Narzulla qizi
1st year student, Faculty of Medicine, Termiz University of Economics and Service
marjonaxusanova489@gmail.com
Ochildiyev Hasan Bahodir o'g'li
Assistant, Termiz State University of Engineering and Agrotechnologies
h1a2s3a4n5ochildiyev@gmail.com

Abstract: Electrical processes in living organisms constitute a fundamental component of biological activity, playing a crucial role in the transmission of nerve signals, muscle contraction, and the regulation of cardiac rhythm. This article provides an in-depth analysis of bioelectric phenomena at the cellular level, focusing on ion transport, the formation of membrane potential, and the mechanisms of action potential propagation. The study is based on classical and modern electrophysiological theories, particularly the Hodgkin–Huxley model, which explains the ionic basis of nerve impulses. Through theoretical and comparative analysis of scientific literature, it is demonstrated that bioelectric phenomena arise due to ion gradients and the selective permeability of cell membranes. In addition, the practical applications of electrocardiography (ECG), electroencephalography (EEG), and electrical stimulation methods in modern medicine are discussed. The results highlight the significant role of bioelectric processes in physiology and clinical diagnostics.

Key words: bioelectric phenomena, membrane potential, ion channels, action potential, electrophysiology, ECG, EEG.

Аннотация: Электрические процессы в живых организмах являются фундаментальным компонентом биологической активности, играя ключевую роль в передаче нервных сигналов, сокращении мышц и регуляции сердечного ритма. В данной статье представлен углублённый анализ биоэлектрических явлений на клеточном уровне с акцентом на транспорт ионов, формирование мембранного потенциала и механизмы распространения потенциала действия. Исследование основано на классических и современных электрофизиологических теориях, в частности на модели Ходжкина–Хаксли, объясняющей ионную природу нервных импульсов. Путём теоретического и сравнительного анализа научной литературы показано, что биоэлектрические явления возникают благодаря градиентам ионов и избирательной проницаемости клеточных мембран. Кроме того, рассматриваются практические применения электрокардиографии (ЭКГ), электроэнцефалографии (ЭЭГ) и методов электрической стимуляции в современной медицине. Полученные результаты подчёркивают важную роль биоэлектрических процессов в физиологии и клинической диагностике.

Ключевые слова: биоэлектрические явления, мембранный потенциал, ионные каналы, потенциал действия, электрофизиология, ЭКГ, ЭЭГ.

Annotatsiya: Tirik organizmlardagi elektr jarayonlari biologik faoliyatning muhim tarkibiy qismi bo'lib, nerv impulslarining uzatilishi, mushaklarning qisqarishi hamda yurak ritmining boshqarilishida muhim rol o'ynaydi. Ushbu maqolada hujayra darajasidagi bioelektrik hodisalar chuqur tahlil qilinib, ionlar transporti, membrana potensialining hosil bo'lishi va harakat potensialining tarqalish mexanizmlariga alohida e'tibor qaratilgan. Tadqiqot klassik va zamonaviy elektrofiziologik nazariyalarga, xususan nerv impulslarining ion asosini tushuntiruvchi Xodjkin–

Xaksli modeliga asoslangan. Ilmiy adabiyotlarning nazariy va taqqosiy tahlili orqali bioelektrik hodisalar ion gradientlari va hujayra membranalarining selektiv o‘tkazuvchanligi natijasida yuzaga kelishi ko‘rsatib berilgan. Shuningdek, zamonaviy tibbiyotda elektrokardiografiya (EKG), elektroensefalografiya (EEG) va elektr stimulyatsiya usullarining amaliy qo‘llanilishi ham ko‘rib chiqilgan. Natijalar bioelektrik jarayonlarning fiziologiya va klinik diagnostikadagi muhim ahamiyatini tasdiqlaydi.

Kalit so‘zlar: bioelektrik hodisalar, membrana potensiali, ion kanallari, harakat potensiali, elektrofiziologiya, EKG, EEG.

Introduction

Electrical activity is an intrinsic property of living organisms and plays a central role in maintaining vital physiological functions. From the transmission of nerve impulses to the rhythmic contraction of the heart, bioelectric processes are essential for the coordination of complex biological systems. At the cellular level, electrical phenomena arise due to the unequal distribution of ions across the cell membrane. This imbalance generates a voltage difference known as the membrane potential, which is crucial for signal transmission in excitable cells such as neurons and muscle fibers.

The scientific study of bioelectricity began with the experiments of Luigi Galvani in the 18th century and was significantly advanced by Hodgkin and Huxley in 1952. Their work provided a quantitative explanation of action potentials based on ion movement, forming the foundation of modern electrophysiology.

Today, bioelectric processes are widely applied in medicine, particularly in diagnostic techniques such as ECG and EEG, making this field highly relevant for both theoretical and applied sciences.

Literature Review

The study of electrical phenomena in biological systems has evolved significantly over time. Early experiments by Galvani demonstrated that muscles respond to electrical stimulation, introducing the concept of “animal electricity.” This idea was further developed through the work of Du Bois-Reymond and Bernstein, who explored membrane potentials and ionic theory. A major breakthrough occurred with the work of Hodgkin and Huxley (1952), who conducted experiments on the squid giant axon and developed a mathematical model describing the ionic mechanisms of action potentials. Their findings showed that sodium (Na^+) influx and potassium (K^+) efflux are responsible for depolarization and repolarization phases.

Recent studies have expanded on this model by investigating ion channels at the molecular level and their role in disease development. Modern electrophysiological research also emphasizes the importance of bioelectric signals in tissue regeneration, neural engineering, and medical diagnostics.

Research Methodology

This research is based on a qualitative, theoretical, and analytical approach aimed at providing a comprehensive understanding of bioelectric processes in living organisms. The study does not involve direct experimental investigation; instead, it relies on the systematic examination and synthesis of existing scientific knowledge in the fields of biophysics, physiology, and electrophysiology. The following research methods were applied:

Analysis of classical and modern scientific literature:

Foundational works, including the studies of Hodgkin and Huxley (1952), as well as contemporary textbooks and peer-reviewed publications, were critically analyzed to identify key theoretical principles and scientific developments related to bioelectric phenomena.

Comparative evaluation of electrophysiological models:

Different theoretical frameworks describing membrane potential and action potential mechanisms were compared to highlight their similarities, differences, and scientific relevance.

Particular emphasis was placed on the explanatory power and limitations of the Hodgkin–Huxley model in comparison with modern interpretations.

Conceptual modeling of ion transport and membrane processes:

The study employed conceptual models to explain the movement of ions (Na^+, K^+, Ca^{2+}) across selectively permeable membranes and their role in generating electrical signals. These models help to simplify complex biological mechanisms and make them analytically accessible.

Inductive generalization and synthesis:

Based on the reviewed literature and theoretical analysis, general conclusions were formulated regarding the nature, origin, and significance of bioelectric processes in living systems.

In addition, an interdisciplinary perspective was adopted, integrating insights from biology, physics, and medical science to ensure a holistic understanding of the topic. The methodological approach emphasizes logical consistency, scientific reliability, and theoretical depth, making the study relevant for both academic research and practical applications in medicine.

Analysis and Results

Membrane Potential Formation

The membrane potential is generated by the unequal distribution of ions across the cell membrane, combined with the selective permeability of the lipid bilayer. Potassium ions (K^+) are present in higher concentrations inside the cell, whereas sodium ions (Na^+) are more abundant in the extracellular fluid. This ionic imbalance is actively maintained by the Na^+/K^+-ATPase pump, which transports three Na^+ ions out of the cell and two K^+ ions into the cell, contributing to the establishment of an electrochemical gradient.

As a result, a stable electrical potential difference is formed, typically around −70 mV in resting neurons. This resting membrane potential is essential for maintaining cellular excitability and readiness for signal transmission. From a biophysical perspective, the membrane behaves as an electrical capacitor, storing charge and enabling rapid voltage changes in response to external stimuli. The analysis of membrane potential demonstrates that even small changes in ion permeability can significantly alter cellular electrical behavior, highlighting the sensitivity of biological systems to ionic dynamics.

Action Potential Mechanism

According to the Hodgkin–Huxley model, the action potential is a rapid, transient change in membrane potential that propagates along excitable cells such as neurons and muscle fibers. This process occurs in several well-defined phases:

Depolarization: Upon reaching a threshold level (approximately −55 mV), voltage-gated sodium channels open, allowing a rapid influx of Na^+ ions into the cell. This causes a sharp rise in membrane potential, often reaching positive values (around +30 mV).

Repolarization: Shortly after depolarization, sodium channels inactivate, and voltage-gated potassium channels open. The efflux of K^+ ions restore the negative membrane potential.

Hyperpolarization: The continued outflow of K^+ ions temporarily increase membrane negativity beyond the resting level, creating a refractory period during which the cell is less excitable.

This sequence follows the “all-or-none” principle, meaning that once the threshold is reached, the action potential is generated with a constant amplitude regardless of stimulus strength. From an analytical standpoint, this mechanism ensures reliable and rapid signal transmission over long distances without loss of intensity. It also prevents signal overlap due to the refractory period, maintaining directional propagation along nerve fibers.

Practical Applications

The understanding of bioelectric mechanisms has led to numerous practical applications in medicine and biomedical engineering. One of the most significant examples is **electrocardiography (ECG)**, which records the electrical activity of the heart and allows clinicians to detect arrhythmias, ischemia, and other cardiac abnormalities. The characteristic P, QRS, and T

waves observed in ECG signals directly reflect the underlying electrical processes of depolarization and repolarization in cardiac tissue.

Another important application is **electroencephalography (EEG)**, which measures the electrical activity of the brain. EEG signals are widely used in the diagnosis of neurological disorders such as epilepsy, sleep disorders, and brain injuries.

In addition, medical devices such as **defibrillators** utilize controlled electrical shocks to restore normal heart rhythm during life-threatening conditions like ventricular fibrillation. Similarly, **neural stimulation technologies** are applied in pain management, rehabilitation, and treatment of neurological disorders.

Results Interpretations

The analysis confirms that bioelectric processes are fundamental to the functioning of living organisms. Electrical signals serve as efficient carriers of information, enabling rapid communication between different organs and systems.

Conclusion

This study confirms that electrical processes in living organisms are essential for maintaining vital physiological functions. Bioelectric phenomena originate from ion gradients across the cell membrane and its selective permeability, which together generate membrane potentials and enable the propagation of action potentials. These processes form the biological basis of nerve impulse transmission, muscle contraction, and cardiac activity.

The analysis shows that bioelectricity is a fundamental physiological mechanism that ensures rapid and coordinated communication within the human body. Electrophysiological models, particularly the Hodgkin–Huxley model, provide a reliable theoretical framework for understanding these processes and remain highly relevant in modern biomedical science.

In addition, the study highlights the practical significance of bioelectric signals in medical diagnostics and treatment. Techniques such as electrocardiography (ECG) and electroencephalography (EEG) allow the non-invasive monitoring of heart and brain activity, demonstrating the applied value of bioelectric research.

Recommendations

1. Further research should focus on the molecular mechanisms of ion channels and membrane transport systems.
2. The development of advanced and more accurate diagnostic technologies based on bioelectric signal analysis is recommended.
3. Bioelectricity and electrophysiology should be more deeply integrated into medical education programs.
4. Expansion of clinical use of electrical stimulation methods in neurology, cardiology, and rehabilitation is encouraged.

References

1. Hodgkin A.L., Huxley A.F. (1952). *A quantitative description of membrane current and its application to conduction and excitation in nerve*. The Journal of Physiology, 116(4), 500–544. Cambridge University Press.
2. Guyton A.C., Hall J.E. (2021). *Textbook of Medical Physiology*. 14th ed. Elsevier, Philadelphia.
3. Bear M.F., Connors B.W., Paradiso M.A. (2020). *Neuroscience: Exploring the Brain*. 4th ed. Lippincott Williams & Wilkins.
4. Alberts B. et al. (2015). *Molecular Biology of the Cell*. 6th ed. Garland Science.
5. Ganong W.F. (2018). *Review of Medical Physiology*. 26th ed. McGraw-Hill Education.
6. Tursunov X.T. (2021). *Biofizika asoslari*. Toshkent: Fan nashriyoti.
7. O‘zbekiston Respublikasi Oliy va o‘rta maxsus ta’lim vazirligi (2022). *Biologiya va biofizika asoslari*. Toshkent.